



# Vector Space

Department of Computer Engineering  
Sharif University of Technology

Hamid R. Rabiee [rabiee@sharif.edu](mailto:rabiee@sharif.edu)

Maryam Ramezani [maryam.ramezani@sharif.edu](mailto:maryam.ramezani@sharif.edu)



# Table of contents

**01**

Review

**02**

Field

**03**

Vector Space

**04**

Linear Combination

**05**

Span-Linear Hull



01

# Review



# Complex Numbers

Sum and Products



# Tuple and Vector Space

## Definition

- A tuple is an ordered list of numbers.
- For example:  $\begin{bmatrix} 1 \\ 2 \\ 32 \\ 10 \end{bmatrix}$  is a 4-tuple (a tuple with 4 elements).

$$\mathbb{R}^2 = \left\{ \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 0.112 \\ \frac{2}{3} \end{pmatrix}, \begin{pmatrix} \pi \\ e \end{pmatrix}, \dots \right\}$$

$$\mathbb{R}^3 = \left\{ \begin{pmatrix} 17 \\ \pi \\ 2 \end{pmatrix}, \begin{pmatrix} 9 \\ -2 \\ \sqrt{2} \end{pmatrix}, \begin{pmatrix} 1 \\ 22 \\ 2 \end{pmatrix}, \dots \right\}$$



# Review: Complex Numbers

Numbers:

- Real: Nearly any number you can think of is a Real Number!

1	12.38	-0.8625	3/4	$\sqrt{2}$	1998
---	-------	---------	-----	------------	------

- Imaginary: When squared give a negative result.

The “unit” imaginary number (like 1 for Real Numbers) is “ $i$ ”, which is the square root of  $-1$ .

Examples of Imaginary Numbers:


$3i$	$1.04i$	$-2.8i$	$3i/4$	$(\sqrt{2})i$	$1998i$
------	---------	---------	--------	---------------	---------

And we keep that little “ $i$ ” there to remind us we need to multiply by  $\sqrt{-1}$


# Review: Complex Numbers

- $\mathbb{C}$  is a plane, where number  $(a + bi)$  has coordinates  $\begin{bmatrix} a \\ b \end{bmatrix}$
- Imaginary number:  $bi$ ,  $b \in \mathbb{R}$

- □ Conjugate of  $x + yi$  is noted by  $\overline{x + yi}$ :
  - $x - yi$



(Complex conjugate)



# Review: Complex Numbers

□ Arithmetic with complex numbers  $(a + bi)$ :

□  $(a + bi) + (c + di) = (a + c) + (b + d)i$

□  $(a + bi)(c + di) = ac - bd + (bc + ad)i$

□  $\frac{a+bi}{c+di} = \frac{(a+bi)(c-di)}{(c+di)(c-di)} = \frac{ac+bd}{c^2+d^2} + \left(\frac{bc-ad}{c^2+d^2}\right)i$




# Review: Complex Numbers

□ Length (magnitude):  $||a + bi||^2 = \overline{(a + bi)}(a + bi) = a^2 + b^2$

• Inner Product:

□ Real:  $\langle x, y \rangle = x_1y_1 + x_2y_2 + \dots + x_ny_n$

□ Complex:  $\langle x, y \rangle = \overline{x_1}y_1 + \overline{x_2}y_2 + \dots + \overline{x_n}y_n$



Extra resource:

If you want to learn more about complex numbers, this video is recommended!

# Binary Operations

What is a binary operation?

# Binary Operations

## Definition

Any function from  $A \times A \rightarrow A$  is a binary operation.

### □ Closure Law:

- A set is said to be closure under an operation (like addition, subtraction, multiplication, etc.) if that operation is performed on elements of that set and result also lies in set.

$$\text{if } a \in A, b \in A \rightarrow a * b \in A$$



# Binary Operations

## Example

- ☐ Is “+” a binary operator on natural numbers?
- ☐ Is “ $\times$ ” a binary operator on natural numbers?
- ☐ Is “-” a binary operator on natural numbers?
- ☐ Is “/” a binary operator on natural numbers?

02

Field



# Fields

$$\forall a, b, c \in F$$

Properties	Binary Operations	
	Addition (+)	Multiplication (.)
Closure (بسته بودن)	$\exists a + b \in F$	$\exists a.b \in F$
Associative (شرکت پذیری)	$a + (b + c) = (a + b) + c$	$a.(b.c) = (a.b).c$
Commutative (جابہ جایی پذیری)	$a + b = b + a$	$a.b = b.a$
Existence of identity $e \in F$	$a + e = a = e + a$	$a.e = a = e.a$
Existence of inverse: For each $a$ in $F$ there <u>must exist</u> $b$ in $F$	$a + b = e = b + a$	$a.b = e = b.a$ <i>For any nonzero <math>a</math></i>
Multiplication is distributive over addition $a.(b + c) = a.b + a.c$ $(a + b).c = a.c + b.c$		

# Fields

- A field in mathematics is a set of things or elements (not necessarily numbers) for which the basic arithmetic operations (addition, subtraction, multiplication, division) are defined:  $(F, +, \cdot)$



## Example


$(\mathbb{R}; +, \cdot)$  and  $(\mathbb{Q}; +, \cdot)$  serve as examples of fields.

- Field is a set  $(F)$  with two binary operations  $(+ , \cdot)$  satisfying following properties:
- 
- 

# Fields


## Example

Set  $B = \{0,1\}$  under following operations is a field?



+	0	1
0	0	1
1	1	0

.	0	1
0	0	0
1	0	1





# Fields

## Example

Which are fields? (two binary operations  $+$ ,  $*$ )

$\mathbb{R}$

$\mathbb{C}$


$\mathbb{Q}$

$\mathbb{Z}$

$\mathbb{W}$

$\mathbb{N}$

$\mathbb{R}^{2 \times 2}$



$\mathbb{C}$  : Complex

$\mathbb{R}$  : Real

$\mathbb{Q}$  : Rational

$\mathbb{Z}$  : Integer

$\mathbb{W}$  : Whole

$\mathbb{N}$  : Natural

03

# Vector Space



# Vector Space

- Building blocks of linear algebra.
- A non-empty set  $V$  with field  $F$  (most of time  $\mathbf{R}$  or  $\mathbf{C}$ ) forms a vector space with two operations:
  1.  $+$  : Binary operation on  $V$  which is  $V \times V \rightarrow V$
  2.  $\cdot$  :  $F \times V \rightarrow V$

## Note

In our course, by **default**, field is  $\mathbf{R}$  (real numbers).

# Vector Space

## Definition

A vector space over a field  $F$  is the set  $V$  equipped with two operations:  $(V, F, +, \cdot)$

- i. **Vector addition:** denoted by “+” adds two elements  $x, y \in V$  to produce another element  $x + y \in V$
- ii. **Scalar multiplication:** denoted by “.” multiplies a vector  $x \in V$  with a scalar  $\alpha \in F$  to produce another vector  $\alpha \cdot x \in V$ . We usually omit the “.” and simply write this vector as  $\alpha x$ .

# Vector Space Properties

## □ Addition of vector space ( $x + y$ )

□ **Commutative**  $x + y = y + x \quad \forall x, y \in V$

□ **Associative**  $(x + y) + z = x + (y + z) \quad \forall x, y, z \in V$

□ **Additive identity**  $\exists \mathbf{0} \in V$  such that  $x + \mathbf{0} = x, \forall x \in V$

□ **Additive inverse**  $\exists (-x) \in V$  such that  $x + (-x) = \mathbf{0}, \forall x \in V$

# Vector Space Properties

## □ Action of the scalars field on the vector space

( $\alpha x$ )

□ **Associative**       $\alpha(\beta x) = (\alpha\beta)x$        $\forall \alpha, \beta \in F; \forall x \in V$

□ **Distributive over** .....

scalar addition:       $(\alpha + \beta)x = \alpha x + \beta x$        $\forall \alpha, \beta \in F; \forall x$

$\in V$

vector addition:       $\alpha(x + y) = \alpha x + \alpha y$        $\forall \alpha \in F; \forall x, y \in V$

□ **Scalar identity**       $1x = x$        $\forall x \in V$

# Vector Space

## Theorem

Every vector space has a unique additive identity.

Every  $v \in V$  has a unique additive inverse.



# Vector Space

## Example

Let  $V$  be the set of all real numbers with the operations  $u \oplus v = u - v$  ( $\oplus$  is an ordinary subtraction) and  $c \odot u = cu$  ( $\odot$  is an ordinary multiplication).

Is  $V$  a vector space? If it's not, which properties fail to hold?



# Vector Space of functions

- Function addition and scalar multiplication

$$(f + g)(x) = f(x) + g(x) \quad \text{and} \quad (af)(x) = af(x)$$


Non-empty set  $X$  and any field  $F$   $\longrightarrow F^X = \{f: X \rightarrow F\}$

## Example

- Set of all polynomials with real coefficients
- Set of all real-valued continuous function on  $[0,1]$
- Set of all real-valued function that are differentiable on  $[0,1]$

# Vector Space of polynomials

$P_n(\mathbb{R})$ : Polynomials with max degree (n)

- 
- Vector addition
  - Scalar multiplication
  - And other 8 properties!



# Vector Space

## Example

Which are vector spaces with  $+$ ,  $*$ ?

- ☐ Set  $\mathbb{R}^n$  over  $\mathbb{R}$
- ☐ Set  $\mathbb{C}$  over  $\mathbb{R}$
- ☐ Set  $\mathbb{R}$  over  $\mathbb{C}$
- ☐ Set  $\mathbb{Z}$  over  $\mathbb{R}$
- ☐ Set of all polynomials with coefficient from  $\mathbb{R}$  over  $\mathbb{R}$
- ☐ Set of all polynomials of degree at most  $n$  with coefficient from  $\mathbb{R}$  over  $\mathbb{R}$
- ☐ Matrix:  $M_{m,n}(\mathbb{R})$  over  $\mathbb{R}$
- ☐ Function:  $f(x): x \rightarrow \mathbb{R}$  over  $\mathbb{R}$

# Conclusion

The operations on field  $F$  are:

- $+: F \times F \rightarrow F$
- $\times: F \times F \rightarrow F$



The operations on a vector space  $V$  over a field  $F$  are:

- $+: V \times V \rightarrow V$
- $\cdot: F \times V \rightarrow V$



04

# Linear Combination



# Linear Combinations

- The **linear combinations** of  $m$  vectors  $a_1, \dots, a_m$ , each with size  $n$  is:

$$\beta_1 a_1 + \dots + \beta_m a_m$$

where  $\beta_1, \dots, \beta_m$  are scalars and called the **coefficients of the linear combination**


- **Coordinates**: We can write any  $n$ -vector  $b$  as a **linear combination of the standard unit vectors**, as:

$$b = b_1 e_1 + \dots + b_n e_n$$

- Example: What are the coefficients and combination for this vector?

$$\begin{bmatrix} -1 \\ 3 \\ 5 \end{bmatrix}$$

# Linear Combinations



Left. Two 2-vectors  $a_1$  and  $a_2$ . Right. The linear combination  $b = 0.75a_1 + 1.5a_2$

## Special Linear Combinations

- ❑ Sum of vectors
- ❑ Average of vectors

05

# Span – Linear Hull



# Span or linear hull

## Definition

If  $v_1, v_2, v_3, \dots, v_p$  are in  $\mathbb{R}^n$ , then the set of all linear combinations of  $v_1, v_2, \dots, v_p$  is denoted by  $\text{Span}\{v_1, v_2, \dots, v_p\}$  and is called the **subset of  $\mathbb{R}^n$  spanned (or generated) by  $v_1, v_2, \dots, v_p$ .**


That is,  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the collection of all vectors that can be written in the form:

$$c_1 v_1 + c_2 v_2 + \dots + c_p v_p$$


with  $c_1, c_2, \dots, c_p$  being scalars.

# Span Geometry

$v$  and  $u$  are non-zero vectors in  $\mathbb{R}^3$  where  $v$  is not a multiple of  $u$




$\text{Span}\{v\}$  as a line through the origin.




$\text{Span}\{u, v\}$  as a plane through the origin.

# Span Geometry



(a)  $\text{Span}(\{\vec{A}, \vec{B}\}) = \text{Span}(\{\vec{A}, \vec{B}, \vec{C}\})$   
 $\text{Span}(\{\vec{A}, \vec{B}, \vec{C}\}) = \text{All vectors on hyperplane}$



(b)  $\text{Span}(\{\vec{A}, \vec{B}\}) \neq \text{Span}(\{\vec{A}, \vec{B}, \vec{C}\})$   
 $\text{Span}(\{\vec{A}, \vec{B}, \vec{C}\}) = \text{All vectors in } \mathbb{R}^3$

Figure 2.6: The span of a set of linearly dependent vectors has lower dimension than the number of vectors in the set

# Span or linear hull

## Example

- ❑ Is vector  $b$  in  $\text{Span} \{v_1, v_2, \dots, v_p\}$
- ❑ Is vector  $v_3$  in  $\text{Span} \{v_1, v_2, \dots, v_p\}$
- ❑ Is vector  $0$  in  $\text{Span} \{v_1, v_2, \dots, v_p\}$
- ❑ Span of polynomials:  $\{(1+x), (1-x), x^2\}$ ?
- ❑ Is  $b$  in  $\text{Span} \{a_1, a_2\}$  ?

$$a_1 = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}, a_2 = \begin{bmatrix} 5 \\ -13 \\ -3 \end{bmatrix}, b = \begin{bmatrix} -3 \\ 8 \\ 1 \end{bmatrix}$$

# Resources

- ❑ Kenneth Hoffman and Ray A. Kunze. Linear Algebra. PHI Learning, 2004.
- ❑ David C. Lay, Steven R. Lay, and Judi J. McDonald. Linear Algebra and Its Applications. Pearson, 2016.
- ❑ Gilbert Strang. Introduction to Linear Algebra. Wellesley-Cambridge Press, 2016.

