



Inner Product Space

Department of Computer Engineering
Sharif University of Technology

Hamid R. Rabiee rabiee@sharif.edu

Maryam Ramezani maryam.ramezani@sharif.edu



Table of contents

01

Introduction

02

Linear Form

03

Bilinear Form

04

**Bilinear Form
on Complex
Vector Space**

05

Inner Product

06


Inner Product Space

01




Introduction



Least Squares Error Correction



Least Squares Error Correction



Error 1: 

Error 2: 

Error 3: 

02

Linear Form



What are Linear Functions?

- $f: R^n \rightarrow R$ means that f is a function that maps real n -vectors to real numbers
- $f(x)$ is the value of function f at x (x is referred to as the argument of the function).
- $f(x) = f(x_1, x_2, \dots, x_n)$: where x_1, x_2, \dots, x_n are arguments

Definition

A function $f: R^n \rightarrow R$ is linear if it satisfies the following two properties:

- **Additivity**: For any n -vector x and y , $f(x + y) = f(x) + f(y)$
- **Homogeneity**: For any n -vector x and any scalar $\alpha \in R$: $f(\alpha x) = \alpha f(x)$

Superposition property:

Definition

Superposition property:

$$f(\alpha x + \beta y) = \alpha f(x) + \beta f(y)$$

Note

❑ A function that satisfies the superposition property is called **linear**

What are Linear Functions?

- If a function f is linear, superposition extends to linear combinations of any number of vectors:

$$f(\alpha_1 x_1 + \cdots + \alpha_k x_k) = \alpha_1 f(x_1) + \cdots + \alpha_k f(x_k)$$

Inner product is Linear Function?

Theorem 1

A function defined as the inner product of its argument with some fixed vector is linear.

Proof?

$$f(x) = a^T x = a_1 x_1 + a_2 x_2 + \cdots + a_n x_n$$

What are Linear Functions?

Theorem 2

If a function is linear, then it can be expressed as the inner product of its argument with some fixed vector.

Proof?

What are Linear Functions?

Theorem 3

The representation of a linear function f in a specific basis as:
 $f(x) = a^T x$ is **unique**, which means that there is only one vector a for which $f(x) = a^T x$ holds for all x .

Proof?

Linear Form Examples

Example

- Is average a linear function?
- Is maximum a linear function?

03

Bilinear Form



Bilinear Form over a real vector space

Definition

Suppose V and W are vector spaces over the same field \mathbb{F} . Then a function $f: V \times W \rightarrow \mathbb{F}$ is called a **bilinear form** if it satisfies the following properties:

a) It is linear in its first argument:

- i. $f(\mathbf{v}_1 + \mathbf{v}_2, \mathbf{w}) = f(\mathbf{v}_1, \mathbf{w}) + f(\mathbf{v}_2, \mathbf{w})$ and
- ii. $f(c\mathbf{v}_1, \mathbf{w}) = cf(\mathbf{v}_1, \mathbf{w})$ for all $c \in \mathbb{F}, \mathbf{v}_1, \mathbf{v}_2 \in V$, and $\mathbf{w} \in W$.

b) It is linear in its second argument:

- i. $f(\mathbf{v}, \mathbf{w}_1 + \mathbf{w}_2) = f(\mathbf{v}, \mathbf{w}_1) + f(\mathbf{v}, \mathbf{w}_2)$ and
- ii. $f(\mathbf{v}, c\mathbf{w}_1) = cf(\mathbf{v}, \mathbf{w}_1)$ for all $c \in \mathbb{F}, \mathbf{v} \in V$, and $\mathbf{w}_1, \mathbf{w}_2 \in W$.

Bilinear Form

Note

Let V be a vector space over a field \mathbb{F} . Then the **dual** of V , denoted by V^* , is the vector space consisting of all linear forms on V .

Example

Let V be a vector space over a field \mathbb{F} . Show that the function $g: V^* \times V \rightarrow \mathbb{F}$ defined by

$$g(f, v) = f(v) \text{ for all } f \in V^*, v \in V$$

is a bilinear form.

Symmetric Bilinear Form

Definition

A **bilinear form** function $f: V \times V \rightarrow \mathbb{F}$ over a real vector space V is called **symmetric** if for all $v, w \in V$:

$$f(v, w) = f(w, v)$$

Bilinear Form arises from a matrix

Theorem 4

Every **bilinear form** function $f: V \times V \rightarrow \mathbb{F}$ over a real vector space V arises from a matrix for all $v, w \in V$:

$$f(v, w) = v^T A w$$

Proof?

Associated Matrices

Definition

If V is a finite-dimensional vector space, $B = \{b_1, \dots, b_n\}$ is a basis of V , and $f: V \times V \rightarrow \mathbb{F}$ be a **bilinear form** function the **associated matrix** A of f with respect to B is the matrix $[f]_B \in \mathbb{F}^{n \times n}$ whose (i, j) -entry is the value $f(b_i, b_j)$.

$$f(v, w) = v^T A w = v^T [f]_B w$$

$$[f]_B = \begin{pmatrix} f(b_1, b_1) & \dots & f(b_1, b_n) \\ \vdots & & \vdots \\ f(b_n, b_1) & \dots & f(b_n, b_n) \end{pmatrix}$$

Associated Matrices

Note

The associated matrix changes if we use a different basis.

Example

For the bilinear form $f\left(\begin{bmatrix} a \\ b \end{bmatrix}, \begin{bmatrix} c \\ d \end{bmatrix}\right) = 2ac + 4ad - bc$ on \mathbb{F}^2 , find $[f]_B$ for basis $B = \left\{\begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 4 \end{bmatrix}\right\}$ and $[f]_P$ for basis $P = \left\{\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right\}$

04

Bilinear Form Over Complex Vector Space



Bilinear Form over a complex vector space

Definition

Suppose V and W are vector spaces over the same field \mathbb{C} . Then a function $f: V \times W \rightarrow \mathbb{C}$ is called a **bilinear form** if it satisfies the following properties:

a) It is **linear in its first argument**:

- i. $f(\mathbf{v}_1 + \mathbf{v}_2, \mathbf{w}) = f(\mathbf{v}_1, \mathbf{w}) + f(\mathbf{v}_2, \mathbf{w})$ and
- ii. $f(\lambda \mathbf{v}_1, \mathbf{w}) = \lambda f(\mathbf{v}_1, \mathbf{w})$ for all $\lambda \in \mathbb{C}, \mathbf{v}_1, \mathbf{v}_2 \in V$, and $\mathbf{w} \in W$.

b) It is **conjugate linear in its second argument**:

- i. $f(\mathbf{v}, \mathbf{w}_1 + \mathbf{w}_2) = f(\mathbf{v}, \mathbf{w}_1) + f(\mathbf{v}, \mathbf{w}_2)$ and
- ii. $f(\mathbf{v}, \lambda \mathbf{w}_1) = \bar{\lambda} f(\mathbf{v}, \mathbf{w}_1)$ for all $\lambda \in \mathbb{C}, \mathbf{v} \in V$, and $\mathbf{w}_1, \mathbf{w}_2 \in W$.

Bilinear Form over a complex vector space



Bilinear forms on \mathbb{R}^n	Bilinear forms on \mathbb{C}^n
<u>Linear</u> in the first variable	<u>Linear</u> in the first variable
<u>Linear</u> in the second variable	<u>Conjugate linear</u> in the second variable



05

Inner Product



Inner product over real vector space

Definition

An inner product is a **positive-definite** **symmetric** **bilinear** form.

An inner product on V is a function $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ such that $v, w \in V, c \in \mathbb{R}$:

1. $\langle v, v \rangle = 0$ if and only if $v = 0$.
2. $\langle w, v \rangle = \langle v, w \rangle$.
3. $\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$ for all $u, v, w \in V$.
4. $\langle cw, u \rangle = c\langle w, u \rangle$ for all $u, w \in V$ and $c \in \mathbb{R}$.
5. $\langle v, v \rangle \geq 0$ for all $v \in V$.

Inner Product

Why for bilinear form I wrote just two properties instead of four properties?

- Using properties (2) and (4) and again (2)

$$\langle w, cu \rangle = \langle cu, w \rangle = c\langle u, w \rangle = c\langle w, u \rangle$$

- Using properties (2), (3) and again (2)

$$\langle w, u + v \rangle = \langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle = \langle w, u \rangle + \langle w, v \rangle$$

1. $\langle v, v \rangle = 0$ if and only if $v = 0$.
2. $\langle w, v \rangle = \langle v, w \rangle$.
3. $\langle u + v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$ for all $u, v, w \in V$.
4. $\langle cw, u \rangle = c\langle w, u \rangle$ for all $u, w \in V$ and $c \in \mathbb{R}$.
5. $\langle v, v \rangle \geq 0$ for all $v \in V$.

Inner Products

Note

□ For $v \in V$, $\langle 0, v \rangle = 0$, $\langle v, 0 \rangle = 0$.



General Inner product

Definition

Suppose that $\mathbb{F} = \mathbb{R}$ or $\mathbb{F} = \mathbb{C}$ and that V is a vector space over \mathbb{F} . Then an **inner product** on V is a function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ such that the following three properties hold for all $c \in \mathbb{F}$ and all $\mathbf{v}, \mathbf{w}, \mathbf{x} \in V$:

- a) $\langle \mathbf{v}, \mathbf{w} \rangle = \overline{\langle \mathbf{w}, \mathbf{v} \rangle}$ (conjugate symmetry)
- b) $\langle \mathbf{v} + c\mathbf{x}, \mathbf{w} \rangle = \langle \mathbf{v}, \mathbf{w} \rangle + c\langle \mathbf{x}, \mathbf{w} \rangle$ (linearity)
- c) $\langle \mathbf{v}, \mathbf{v} \rangle \geq 0$, with equality if and only if $\mathbf{v} = \mathbf{0}$. (pos. definiteness)

Inner Products for vectors

Note

- The standard inner product between vectors is: $(x, y \in \mathbb{R}^n)$

$$\langle x, y \rangle = x^T y = \sum x_i y_i$$

- The function $\langle \cdot, \cdot \rangle : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}$ defined by

$$\langle v, w \rangle = v^* w = \sum_{i=1}^n \bar{v}_i w_i$$

for all $v, w \in \mathbb{C}^n$ is an inner product on \mathbb{C}^n .

Inner Products for matrices

Note

The standard inner product between two matrices is: $(X, Y \in \mathbb{R}^{m \times n})$

$$\langle X, Y \rangle = \text{trace}(X^T Y) = \sum_i \sum_j X_{ij} Y_{ij}$$

Example

Find the inner products of following matrices:

$$U = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}, \quad V = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$$

Inner Product for functions

Note

Let $a < b$ be real numbers and let $C[a, b]$ be the vector space of continuous functions on the real interval $[a, b]$. The function $\langle \cdot, \cdot \rangle : C[a, b] \times C[a, b] \rightarrow \mathbb{R}$ defined by

$$\langle f, g \rangle = \int_a^b f(x)g(x) dx \quad \text{for all } f, g \in C[a, b]$$

is an inner product on $C[a, b]$.

Inner Product for polynomials

Note

□ For $p(x)$ and $q(x)$ with at most degree n :

$$\langle p(x), q(x) \rangle = p(0)q(0) + p(1)q(1) + \cdots + p(n)q(n)$$

□ For $p(x)$ and $q(x)$: $\langle p(x), q(x) \rangle = p(0)q(0) + \int_{-1}^1 p'q'$

□ For $p(x)$ and $q(x)$: $\langle p(x), q(x) \rangle = \int_0^\infty p(x)q(x)e^{-x}dx$

06

Inner Product Space

Inner product space

Definition

An **inner product space** is a finite-dimensional real or complex vector space V along with an inner product on V .

Euclidean Space Unitary Space

Resources

- ❑ Chapter 8: Kenneth Hoffman and Ray A. Kunze. Linear Algebra. PHI Learning, 2004.
- ❑ Chapter 6: Sheldon Axler, Linear Algebra Done Right, 2024.
- ❑ Chapter 1: David C. Lay, Steven R. Lay, and Judi J. McDonald. Linear Algebra and Its Applications. Pearson, 2016.
- ❑ Chapter 2: David Poole, Linear Algebra: A Modern Introduction. Cengage Learning, 2014.
- ❑ Chapter 1: Gilbert Strang. Introduction to Linear Algebra. Wellesley-Cambridge Press, 2016.